


October, 1979
(Purchase Order 40-9AD6-9-811)



Ecological Survey of the Vegetation of the proposed
Smoky Creek Research Natural Area,
Shasta-Trinity National Forest, Trinity County, California

Report to the R5/PSW Research Natural Area
Committee, U.S. Forest Service

D.W. Taylor and K.A. Teare
Botany Department
University of California
Davis, CA 95616

| <u>Table of Contents</u> | <u>Page</u> |
|--|-------------|
| Abstract..... | 2 |
| Introduction..... | 2 |
| Location..... | 2 |
| Physiography..... | 2 |
| Regional Climate..... | 3 |
| Geology and Soils..... | 4 |
| Vegetation..... | 5 |
| Regional Vegetation..... | 5 |
| Sampling Methods..... | 6 |
| Plant Associations..... | 6 |
| Forest Cover Types..... | 12 |
| Conclusions and Recommendations..... | 12 |
| Plant Species List..... | 16 |
| Regional Map..... | 27 |
| Location Map..... | 28 |
| Topographic Map..... | 29 |
| Vegetation Map..... | 30 |
| Climatic Summary..... | 31 |
| Riparian, Seep and Meadow Vegetation association Table..... | 32 |
| Forest vegetation association Table..... | 33 |
| Forest Plot Summary table..... | 34 |
| Appendix 1-Forest Plot Data..... | 36 |
| Appendix 2-Formal nomenclature of vegetation associations..... | 40 |

Abstract

This report details the results of an ecological survey of the vegetation of the proposed Smoky Creek Research Natural Area. The area is being considered for reclassification to Research Natural Area status by the Forest Service, U.S. Department of Agriculture. This report describes the ecological setting of the proposed Research Natural Area in relation to local gradients in vegetation composition.

INTRODUCTION

Location. The proposed Smoky Creek Research Natural Area is located on several minor tributary streams of the South Fork of the Trinity River about 5 miles SE from Forest Glen, Trinity County, California ($40^{\circ}22'N, 125^{\circ}15'W$, See Maps 1 & 2). The proposed Research Natural Area boundaries circuitously skirt nearby timbersales at the headwaters of Smoky Creek and Silver Creek in portions of Sections 20, 21, 28, and 29 of T27N, R12W, M.D.M. The area boundaries encompass 250 ha as they are drawn by the Shasta-Trinity National Forest (see Map 3).

Physiography. The Smoky Creek Research Natural Area lies at the head of the easternmost fork of Silver Creek and on the North Fork of Smoky Creek. The area is physiographically a portion of the canyon of the South Fork of the Trinity River, and is near a summit separating the headwaters of the latter stream from those which drain to the Sacramento Valley to the east. Elevations on the proposed Research Natural Area range from about 1000 m (3200 ft) to about 1300 m (4200 ft).

Slopes within the boundaries of the area are generally on the order of 10° - 20° , and there are extensive, broad interfluvies of lesser slope between Smoky Creek and Silver Creek.

Both Silver Creek and the North Fork of Smoky Creek are ephemeral streams. At the time of our visit (September 1979), they held water only in isolated sections, and are probably completely dry by late-summer in years with less than normal precipitation. The springs in nearby Swim Meadow and at Silver Spring flow throughout the summer months, as do several local unmapped seeps.

The proposed Smoky Creek Research Natural Area is easily accessible via Forest Road 29 (Bramlet Road) off Highway 36 about 7 miles to the east of Forest Glen. The Smoky Creek Trail (12W35) shown on the 1969 Shasta-Trinity National Forest Visitor's Map is signed at its junction with Forest Road 29. This trail is passable by four-wheel drive vehicle into approximately the center of the Research Natural Area. Other trails shown on the Forest Visitor's Map are overgrown and are hard to follow or to detect on aerial photography.

Regional Climate. The nearest recording station in the vicinity of the proposed Research Natural Area is at Forest Glen, 5 miles to the NW and nearly 450 m lower in altitude. The climatic data for Forest Glen are summarized in Table 1. The actual climate of the Smoky Creek area is probably somewhat cooler and wetter than at Forest Glen, owing to higher altitude. The mean annual temperature at Forest Glen is 10.8°C (51°F), and the mean annual precipitation is 1572 mm. Table 1 also gives

an estimated mean monthly water balance (Major 1977) using the Forest Glen data and assuming 100 mm of soil water storage capacity. Potential Evapotranspiration (PE in Table 1) is estimated to be 664 mm/yr, while Actual Evapotranspiration (AE in Table 1) is estimated at 394 mm/yr. Forest growth is thus probably limited on the most severe topoedaphic sites in this region by water stress. The area probably receives moderate snowfalls in winter.

Geology and Soils. The proposed Smoky Creek Research Natural Area is underlain by Pre-Cretaceous metamorphic rocks, with extensive outcroppings of altered ultrabasic intrusive rock of the same relative age-serpentine. The regional geology has been mapped at a scale of 1:125,000 (Cal. Div. Mines & Geol. 1962), with reconnaissance geology at a scale of 1:24,000 from the South Fork of the Trinity River to the northeast (Hicks 1960).

Soils in the vicinity of Smoky Creek have not been mapped in published form at the series level, nor are soil-vegetation maps available (Colwell 1977). Published soil descriptions for the vicinity of the proposed Research Natural Area are available for Tehama County just to the east (Gowan et al. 1967)

The Sheetiron-Josephine soil association described by Gowan et al. develop over hard sedimentary rocks, and are moderately deep, steep or very steep stony soils. The Dubakella-Neuns soils are moderately deep or deep, steep or very steep, stony soils which develop over altered volcanic rocks. Serpentine derived soils of the region are related to this latter complex.

VEGETATION

Regional Vegetation. The proposed Smoky Creek Research Natural Area is located in a region of rapid vegetation transition. Coastward, diverse, mesic coniferous forests are typical, while more inland sites support xeric foothill communities. Sawyer et al. (1977), and Sawyer and Thornburgh (1977) review the diverse vegetation patterns in this general geographic region, but their coverage is based on a limited literature on vegetation composition. The zonal forests of the region are dominated by Pseudotsuga menziesii. Taylor (1975) described the Pseudotsuga forest vegetation across the canyon of the South Fork of the Trinity River from the Smoky Creek area. At this site, on the generally NE facing slopes at the base of South Fork Mountain, Pseudotsuga was the canopy dominant, with minor amounts of Pinus lambertiana, Pinus ponderosa and Abies concolor. Arbutus menziesii, Chrysolepis chrysophylla and Calocedrus decurrens were also present in the tree stratum. The understory of these different associations was open to dense, with xeromesic shrubs such as Berberis nervosa. Taylor (1975) named this association the Pseudotsuga menziesii-Berberis nervosa Association.

Serpentine vegetation is also a very important feature of the region. Pinus jeffreyi forms open to semi-dense stands on the more heavily mineralized sites, and a number of rare plant associates occur in these habitats, including Arenaria rosei, Haplopappus ophitidis, Streptanthus barbatus, Asclepias solonana, Eriogonum siskiyouense and Allium hoffmanii (Stebbins 1972).

Survey Methods. We visited the Smoky Creek area on 12-15 September 1979. At that time we sampled 15 plots and 24 releves. Releves were summary lists made in representative habitat types (see Mueller-Dombois and Ellenberg 1974). Each taxon present in a homogeneous area of vegetation was enumerated, and cover of each was visually estimated in the following cover classes: greater than 75%=5; 50-75% cover = 4; 25-50% cover = 3; 5-25% cover = 2; 1-5% cover = 1; less than 1% cover but numerous individuals = +; few individuals of low cover = R. These cover/abundance data form the data entries in Tables 2 and 3.

At selected stands of forest vegetation, 10 m radius circular plots (1 at 15 m radius) were established. Diameter of all stems 1.37 m tall was measured to the nearest .1 cm. A releve was also made at each plot locality.

The releve data was coded and recorded on punch cards for analysis. Association Tables (Tables 2 & 3) were ordered using the algorithm of Ceska and Roemer (1971) as modified by Taylor (1976). Units of vegetation (associations) were determined from this arranged primary data matrix. The stand data were similarly reduced by computer. Appendix 1 gives the summarized data for each plot. Basal Area (m^2/ha), Relative Dominance (Proportion total Basal Area), Density (Stems/ha), Relative Density (Proportion total density), Importance Value (Relative Dominance + Relative Density), and mean stem diameter (cm) were calculated.

Plant Associations. The 39 releves sampled were organized by tabular association analysis (as shown in Tables 2 & 3) to identify groups of similar species with regard to their cooccurrence in the habitats

sampled. From this arranged matrix, 8 associations can be recognized based on the presence/absence of the species groups identified in this analysis. These 8 vegetation types are related in a hierarchical fashion as follows:

- A. Alnus rhombifolia-Peltiphyllum peltatum Alliance
 - 1. Alnus rhombifolia-Galium triflorum Association
 - 2. Prunella vulgaris lanceolata-Carex bolanderi Association
- B. Deschampsia caespitosa Alliance
 - 3. Deschampsia caespitosa-Carex buxbaumii Association
- C. Carex rostrata Alliance
 - 4. Carex exsiccata Association
- D. Pinus jeffreyi-Onychium densum Alliance
 - 5. Pinus jeffreyi-Haplopappus ophitidis Association
 - 6. Ceanothus cuneatus-Sitanion hystrix Association
 - 7. Pinus jeffreyi-Calamagrostis koelerioides Association
- E. Pseudotsuga menziesii Alliance
 - 8. Pseudotsuga menziesii-Pinus ponderosa Association

A. Alnus rhombifolia-Peltiphyllum peltatum Alliance.

This is the major riparian vegetation type along major watercourses in the mountains of California (Knapp 1965). Along the South Fork of the Trinity River below the Smoky Creek area, this vegetation occurs in its nominate compositional form.

1. Alnus rhombifolia-Galium triflorum Association (SAF 221)

This association is the ephemeral stream phase of the A. rhombifolia-P. peltatum alliance. The latter characteristic dominant of the herb layer is absent from Smoky Creek at the upper elevations, owing to low summer flows. A typical stand of this association has a sparse overstory of

Alnus rhombifolia, with Acer macrophyllum and Pseudotsuga as codominants, averaging 11.0, 1.1 and 3.1 m²/ha Basal Area respectively. Mesic shrubs are common in the understory, including Rosa pisocarpa, Philadelphus lewisii gordonianus, Rubus parviflorus and Euonymus occidentalis. Galium triflorum is a mesic site herb which dominates the sparse herb layer in this association.

2. Prunella vulgaris lanceolata-Carex bolanderi Association

The vegetation typified by this association is characteristic of seeps and small springs in the Smoky Creek area. Some typical Alnus rhombifolia-Galium triflorum species are present, but this vegetation is a floristically distinct type, with a meadow physiognomy. Carex bolanderi is common, and C. amplifolia is often a codominant along with Torreyochloa pauciflora. Prunella vulgaris lanceolata is a constant species, and also occurs in the previously described association.

C. Deschampsia caespitosa Alliance.

This alliance is an important meadow vegetation unit in the mountains of the Holoarctic (cf. Komarkova 1979). Deschampsia caespitosa is a circumboreal species which dominates these wet, high productivity meadows.

3. Deschampsia caespitosa-Carex buxbaumii Association

This vegetation occurs in the Silver Spring and Swim Meadow areas adjacent to the proposed Smoky Creek Research Natural Area. These meadows are supplied by year-round flow of highly mineralized water, and they support a distinctive and uncommon vegetation. The mineralization of these seeps is likely to be unusual due to the close proximity of large

expanses of serpentine nearby. Swim Meadow is dominated by Deschampsia, Juncus covillei and J. effusus. Carex buxbaumii and Helenium bigelovii are conspicuous codominants. Silver Spring appears more heavily mineralized, with the presence of Sisyrinchium idahoense indicating this condition. Rhynchospora glomerata minor is a very rare sedge-like plant which is codominant at Silver Spring. Rhynchospora glomerata var. minor is known from only 6 other localities in California, and the Silver Spring population represents a new collection. Other distinctive taxa occur in these serpentine seeps, including Cirsium breweri and Hastingsia alba.

C. Carex rostrata Alliance.

This vegetation is characteristic of shallow, often ephemeral pools and ponds in the northern hemisphere mountains. Carex taxa of section Vesicarieae are the indicator group-as they can tolerate the oligotrophic, highly acidic flooded conditions.

4. Carex exsiccata Association

One small pond vegetated by C. exsiccata was seen in the Smoky Creek drainage. Carex exsiccata is the low elevation vicariant of C. vesicarya and C. rostrata, which both dominate wet meadows in much of mountainous California.

E. Pinus jeffreyi-Onychium densum Alliance (SAF 247)

This vegetation is confined to serpentine substrates in the California North Coast Ranges. Whittaker (1960) describes the more northern phase of this vegetation in the Siskiyou Mountains. Serpentine is very common at mid-elevations in the Smoky Creek area, so this vegetation is an important landscape element.

5. Pinus jeffreyi-Haplopappus ophitidis Association

A typical stand of this vegetation is very open, with widely spaced but large Pinus jeffreyi averaging 300 stems/ha and $36 \text{ m}^2/\text{ha}$ Basal Area. Calocedrus decurrens is a scattered codominant. The shrub understory cover is sparse, with small Ceanothus cuneatus being most common. This association is characteristic of the rawest serpentine sites, and is differentiated by the presence of the low subshrub Haplopappus ophitidis. Calamagrostis koelerioides is a dominant grass in the herb layer.

6. Ceanothus cuneatus-Sitanion hystrix Association.

This association is found on very rocky soils or serpentine outcrops in the Smoky Creek area. Ceanothus cuneatus forms a dense (about 50% cover) low shrub (1 m tall) canopy. Herbs and occasional associated shrub species which form the understory of the Pinus jeffreyi-Haplopappus ophitidis and P. jeffreyi-Calamagrostis koelerioides are present. Soils are poorly developed in these sites, precluding the establishment of P. jeffreyi, although seedlings or stunted small individuals can occasionally be seen here.

7. Pinus jeffreyi-Calamagrostis koelerioides Association

A typical stand of this vegetation is physiognomically similar to the P. jeffreyi-Haplopappus ophitidis association, in that the taxa which differentiate the latter association are small herbs or subshrubs. The overstory characteristics of the two P. jeffreyi vegetation types are similar, with P. jeffreyi averaging $32.8 \text{ m}^2/\text{ha}$ and 381 stems/ha in this association. This vegetation occurs on serpentine derived soils, but in less heavily mineralized or more highly weathered sites than the P. jeffreyi-H. ophitidis association.

E. Pseudotsuga menziesii Alliance

Pseudotsuga menziesii vegetation is the zonal climatically controlled climax vegetation in the general region of the proposed Smoky Creek Research Natural Area. This vegetation is near to its upper elevational limits in the study area, but is a generally xeric facies of this vegetation at Smoky Creek because of the generally S or W exposures and because of the shallow and partly or totally serpentine derived soils.

8. Pseudotsuga menziesii-Pinus ponderosa Association (SAF 244)

This vegetation is the most conspicuous landscape feature of the proposed Smoky Creek Research Natural Area. It grades into the Pinus jeffreyi-Onychium densum alliance very infrequently, and usually only where significant colluviation has occurred. Pseudotsuga menziesii-Pinus ponderosa vegetation develops only over non-serpentine substrates in the study area. Following disturbance, Arctostaphylos viscida stands become established on the more xeric sites of this habitat type, and Ceanothus integerrimus stands on the more mesic ecotopes. Neither of these two subassociations were sampled as they are of only local importance on the proposed Research Natural Area.

A typical stand of Pseudotsuga menziesii-Pinus ponderosa vegetation is open to semi-dense. The coniferous overstory is generally older age class individuals of low density. Dense to semi-dense conifer reproduction may occur and is typically patchy. Pseudotsuga is the canopy dominant, averaging 49.6 m²/ha Basal Area and 498 stems/ha. Pinus lambertiana and Pinus ponderosa share codominance at 14.6 and 18.2 m²/ha Basal Area respectively, and are both found at about equal densities

of 100 stems/ha. Quercus kelloggii and Q. chrysolepis form a second tree layer. The shrub understory is open, with Toxicodendron diversilobum and an occasional Arctostaphylos viscida or Ceanothus integerrimus. The herb layer is generally sparse, with few constant or dominant species (cf Table 3).

Forest Cover Types. The forest vegetation of the proposed Smoky Creek Research Natural Area typifies the following SAF Forest Cover Types (Soc. American Foresters 1954):

SAF 221 Red Alder (White Alder variant)

SAF 245 Douglas Fir-Ponderosa Pine

SAF 247 Jeffrey Pine

The proposed natural area fully characterizes the spatial and temporal variance of the Douglas Fir-Pacific Ponderosa Pine type in the general vicinity of the headwaters of the South Fork of the Trinity River.

CONCLUSIONS AND RECOMMENDATIONS

Several sensitive plant species occur on the serpentine areas to the east of the proposed Research Natural Area eastern boundary. Map 2 shows some localities of several of these rare plants (data from B. Williams, Shasta-Trinity National Forest). We collected Haplopappus ophitidis on the serpentine in the SW $\frac{1}{4}$ of Section 22, T27N, R12W, but did not observe others of these rare plant taxa given on Map 2. Since our visit was late in the growing season, their presence on the proposed Natural Area can not be ruled out until a full botanical inventory of the region is undertaken.

Rhyncospora glomerata var. minor is a rare sedge-like plant we collected at Silver Spring. It is listed as RARE (R.E.V.D. Code 2-2-2-1, Powell 1974) by the California Native Plant Society. It is only known from six other, widely scattered localities in California.

We also collected Helianthus exilis Gray (R.E.V.D. Code 2-2-?-3) on a vernal seep underlain by serpentine within the proposed Research Natural Area boundaries. Munz (1959) incorrectly followed the treatment of Heiser et. al (1969), and relegated H. exilis to synonymy under H. bolanderi Gray. Heiser (1949) has speculated that H. bolanderi is a very recent derivative taxon which has arisen through introgressive hybridization between H. exilis and the weedy introduced H. annuus L.. Helianthus annuus was introduced to California with the first extensive agriculture, and entered the serpentine belt, according to Heiser's scenario, where H. exilis is native. Our collection of H. exilis some 75 miles north of its previously known range, and the complete lack of H. bolanderi or H. annuus would seem to corroborate Heiser's hypothesis. Heiser's mistake is that H. exilis is the native taxon, and should retain species status, while H. bolanderi is simply a recent hybrid. Thus, a non-introgressed H. exilis population (analagous to a pure strain of any of our Western North American trout not affected by hybridization with introduced strains of rainbow trout) is of critical interest to this impacted and rare species.

The proposed boundaries of the Smoky Creek Research Natural Area are drawn to include the most extensive stands of Pseudotsuga menziesii-Pinus ponderosa vegetation in the drainages of the study area which have

not been disturbed by logging. While this is perfectly valid, it does exclude some very distinct habitats nearby-namely Silver Spring and the heavily serpentized ridge just to the east of the easternmost boundary. We recommend the proposed boundaries be adjusted to include these nearby features. It is also our understanding that the Shasta-Trinity National Forest is planning to contract out for a botanical survey in 1980 of the serpentine Pinus jeffreyi areas in the vicinity of the proposed Smoky Creek Research Natural Area. We would recommend that such a survey consider the proper easternmost boundaries if it is felt that inclusion of additional P. jeffreyi vegetation within the proposed Research Natural Area is warranted.

Literature Cited

- Ceska, A. and H. Roemer. 1971. A computer program for identifying species releve groups in vegetation studies. *Vegetatio* 23:255-277.
- Colwell, W. L., Jr. 1977. The status of vegetation mapping in California. Chapter 6 in M.G. Barbour and J. Major (Eds.) *Terrestrial Vegetation of California*. Wiley, Interscience, New York. pp. 195-222.
- California Division of Mines and Geology. 1962. Redding sheet. Compiled by Rudolph G. Strand.
- Gowan, et. al. 1967. Soil survey of Tehama County, California. U.S.D.A., Soil Conservation Service.
- Heiser, C.B., Jr., D.M. Smith, S.B. Clevenger and W. C. Martin, Jr. 1969. The North American Sunflowers (*Helianthus*). *Mem. Torrey Bot. Club* 22:1-218
- _____. 1949. Study in the evolution of the sunflower species *Helianthus annuus* and *H. bolanderi*. *Univ. Calif. Pubs. Bot.* 23: 157-196.
- Hicks, B. 1960. Geology of the Burnt Ranch Dam Site, Trinity County, California. Calif. Div. Mines and Geol. unpublished open file report.

- Knapp, R. 1965. Die vegetation von Nord- und Mittelamerika und der Hawaii-Inslan. Fischer, Stuttgart.
- Major, J. 1977. California climate in relation to vegetation. Ch. 2 in M.G. Barbour and J. Major (Eds.) Terrestrial Vegetation of California. Wiley, Interscience, New York. pp. 11-74.
- Müller-Dombois, D., and H. Ellenberg. 1974. Aims and methods of vegetation ecology. Wiley, Interscience, New York.
- Society of American Foresters. 1964. Forest Cover types of North America.
- Sawyer, J.O., D. A. Thornburgh and J. R. Griffin. 1977. Mixed evergreen forest. Ch. 10 in M.G. Barbour and J. Major (Eds.) Terrestrial Vegetation of California. Wiley, Interscience, New York.
- _____, and _____. 1977. Montane and subalpine vegetation of the Klamath Mountains. Ch. 20 in M.G. Barbour and J. Major (Eds.) Terrestrial Vegetation of California. Wiley, Interscience, New York.
- Stebbins, G. L. 1972. The Yolla Bolly serpentines. Calif. Native Plant Soc. Bull. Vol. 8(3):5-7.
- Taylor, D. W. 1975. Structure of an old-growth Douglas-Fir forest in northwestern California. Unpublished PSW RNA Establishment Report.
- _____. 1976. Ecology of the timberline vegetation at Carson Pass, Alpine County, California. Ph.D. Dissertation, Univ. California, Davis.
- Whittaker, R. H. 1960. Vegetation of the Siskiyou mountains, Oregon and California. Ecol. Monographs 30:279-338.

Plant species list for the proposed Smoky Creek Research Natural Area, Shasta-Trinity National Forest, Trinity County, California. The number to the right of each taxon is the page number where the taxon can be referenced in Munz and Keck's A California Flora, Munz' Supplement to A California Flora (indicated by S), and Munz's A Flora of Southern California (indicated by FSC).

ACERACEAE

Acer macrophyllum Pursh 996
Riparian. Common.

AMARYLLIDACEAE

Allium sp. 1368
Rocky slopes; serpentine outcrops. Uncommon.
Triteleia laxa Benth. 1381
Rocky slopes. Uncommon.

ANACARDIACEAE

Rhus trilobata Nutt. ex T. & G. var. quinata Jeps. 998
Riparian; ephemeral streambeds. Uncommon.
Toxicodendron diversilobum (T. & G.) Greene 998, 67 FSC
Riparian; mixed conifer forest; rocky talus slopes. Common.

APOCYNACEAE

Apocynum pumilum (Gray) Greene 451
Mixed conifer forest. Frequent.

ARALIACEAE

Aralia californica Wats. 1000
Riparian. Frequent.

ARISTOLOCHIACEAE

Asarum hartwegii Wats. 965
Mixed conifer forest. Infrequent.

ASPIDIACEAE

Polystichum munitum (Kaulf.) Presl. 40
var. imbricans (D.C. Eat.) Maxon
Rocky talus slopes. Frequent.

BETULACEAE

- Alnus rhombifolia Nutt. 900
Riparian; seeps. Frequent.
Corylus cornuta Marsh. var. californica (A. DC.) Sharp. 899
Riparian; mixed conifer forest. Common.

BLECHNACEAE

- Woodwardia fimbriata Sm. in Rees 44
Seeps in mixed conifer forest. Infrequent.

CAMPANULACEAE

- Campanula prenanthoides Durand. 1063
Mixed conifer forest. Infrequent.

CAPRIFOLIACEAE

- Lonicera hispidula Dougl. var. vacillans Gray 1051
Seeps. Infrequent.
Symphoricarpos mollis Nutt. in T. & G. 1049
Seeps in mixed conifer forest.

CARYOPHYLLACEAE

- Cerastium arvense L. 277
Seep. Infrequent.
Silene californica Durand 288
Mixed conifer forest. Uncommon.

CELASTRACEAE

- Euonymus occidentalis Nutt. ex Torr. 967
Riparian. Uncommon.

COMPOSITAE

- Achillea lanulosa Nutt. 1228
Serpentine outcrop. Common.
Adenocaulon bicolor Hook 1239
Mixed conifer forest. Common.
Anaphalis margaritacea (L.) Benth. ex Clarke 1263
Common along bars of dry streambeds.
Arnica cordifolia Hook 1243
Mixed conifer forest. Common.
Artemisia douglasiana Bess. in Hook 1237
Serpentine outcrop. Common.
Aster adscendens Lindl. in Hook 1199
Swim Meadow. Infrequent.
Balsamorhiza sagittata (Pursh) Nutt. 1086
Serpentine outcrops. Frequent.

COMPOSITAE

| | |
|---|------|
| <u>Chrysothamnus parryi</u> (Gray) Greene | 1189 |
| ssp. <u>latior</u> Hall & Clem. | |
| Serpentine outcrops. Frequent. | |
| <u>Cirsium arvense</u> (L.) Scop. | 1280 |
| Seeps. Introduced. Frequent. | |
| <u>Cirsium breweri</u> (Gray) Jeps. | 1278 |
| Serpentine outcrops; Swim Meadow. Frequent. | |
| <u>Cirsium callilepis</u> (Greene) Jeps. | 1277 |
| Serpentine outcrops. Frequent. | |
| <u>Eriophyllum lanatum</u> (Pursh) Forbes | 1146 |
| var. <u>lanceolatum</u> (Howell) Jeps. | |
| Serpentine outcrops. Frequent. | |
| <u>Eupatorium occidentale</u> Hook | 1268 |
| Riparian. Common. | |
| <u>Haplopappus ophitidis</u> (J.T. Howell) Keck | 1179 |
| Serpentine patch on divide between Yolla Bolly | |
| and Hayfork ranger districts. Infrequent. | |
| <u>Helenium bigelovii</u> Gray | 1139 |
| Swim Meadow; Silver Spring. Frequent. | |
| <u>Helianthus bolanderi</u> Gray | 1088 |
| cf. <u>Helianthus exilis</u> Gray | |
| Serpentine seeps. Infrequent. | |
| <u>Hieracium albiflorum</u> Hook | 1305 |
| Seeps in mixed conifer forest. Frequent. | |
| <u>Hieracium greenei</u> Gray | 1305 |
| Serpentine outcrops. Infrequent. | |
| <u>Lactuca saligna</u> L. | 1304 |
| Along meadow border at Swim Meadow. Infrequent. | |
| <u>Lessingia nemaclada</u> Greene | 1222 |
| Brushy slopes. Frequent. | |
| <u>Madia elegans</u> D. Don | 1115 |
| Ephemeral streambeds. Frequent. | |
| <u>Petasites palmatus</u> (Ait.) Gray | 1253 |
| Seeps. Infrequent. | |
| <u>Senecio greenei</u> Gray | 1250 |
| Serpentine outcrops. Frequent. | |
| <u>Wyethia angustifolia</u> (DC.) Nutt. | 1085 |
| Mixed conifer forest. Uncommon. | |

CONVOLVULACEAE

| | |
|---------------------------------|-----|
| <u>Convolvulus</u> sp. | 459 |
| Mixed conifer forest. Uncommon. | |

CORNACEAE

| | |
|----------------------------------|------|
| <u>Cornus nuttallii</u> Aud. | 1035 |
| Seeps. Common. | |
| <u>Cornus stolonifera</u> Michx. | 1034 |
| Riparian. Infrequent. | |

CRASSULACEAE

- Sedum obtusatum Gray ssp. boreale Clausen 727
Rocky talus slopes. Infrequent.

CRUCIFERAE

- Erysimum capitatum (Dougl.) Greene 268
Rocky slopes. Infrequent.
Streptanthus barbatus Wats. 218
Serpentine outcrops. Infrequent.
Streptanthus tortuosus Kell. var. pallidus Jeps. 218
Rocky slopes. Infrequent.

CRYPTOGRAMMATACEAE

- Onychium densum Brack. in Wilkes 37
Serpentine outcrops. Common.

CUPRESSACEAE

- Calocedrus decurrens (Torr.) Florin 59, 7S
Mixed conifer forest, serpentine outcrops. Common.

CUSCUTACEAE

- Cuscuta subinclusa Dur. & Hilg. 467
Infrequent.

CYPERACEAE

- Carex amplifolia Boott. 1456
Seeps. Common.
Carex bolanderi Olney 1443
Seeps. Common.
Carex buxbaumii Wahl. 1458
Swim Meadow. Common.
Carex exsiccata Bailey 1462
Swim Meadow; seeps and vernal pool. Infrequent.
Carex globosa Boott. 1450
Mixed conifer forest, serpentine outcrops. Infrequent.
Carex multicaulis Bailey 1451
Riparian. Frequent.
Carex nudata W. Boott. 1460
Riparian. Frequent.
Heleocharis montevidensis Kunth. 1420
var. parishii (Britton) V. Grant
Uncommon in Swim Meadow.
Rhynchospora glomerata (L.) Vahl. var. minor Britton 1428
Silver Spring. Infrequent.

DATISCACEAE

- Datisca glomerata (Presl.) Baill. 177
Riparian; seeps. Infrequent.

DENNSTAEDTIACEAE

- Pteridium aquilinum (L.) Kuhn var. pubescens Underw. 32
Seeps. Frequent.

EQUISETACEAE

- Equisetum arvense L. 28
Silver Spring; seeps. Common.
Equisetum laevigatum A. Br. 27
Seeps. Common.
Equisetum telmateia Ehrh. var. braunii Milde 27
Seeps. Infrequent.

ERICACEAE

- Arbutus menziesii Pursh. 415
Mixed conifer forest. Infrequent.
Arctostaphylos manzanita Parry 422
Serpentine outcrops. Frequent.
Arctostaphylos viscida Parry 424
Mixed conifer forest. Frequent.

FAGACEAE

- Chrysolepis chrysophylla (Dougl. ex Hook) Hjelmquist 901, 120S
Seeps. Infrequent.
Quercus chrysolepis Liebm. 906
Mixed conifer forest. Common.
Quercus durata Jeps. 906
Serpentine outcrops. Frequent.
Quercus garryana Dougl. 904
Mixed conifer forest; serpentine outcrops. Frequent.
Quercus kelloggii Newb. 903
Mixed conifer forest. Common.

GRAMINEAE

- Agrostis exarta Trin. 1522
Vernal pond. Frequent.
Bromus marginatus Nees. 1470
Riparian; seeps; serpentine outcrops. Common.
Bromus tectorum L. 1474
Grassy meadows; brushy slopes. Frequent.
Calamagrostis koelerioides Vasey 1518
Serpentine outcrops. Common.
Deschampsia caespitosa (L.) Beauv. 1513
Swim Meadow; Silver Spring. Common.

GRAMINEAE

| | |
|--|------|
| <u>Elymus glaucus</u> Buckl. | 1505 |
| Riparian; seeps. Common. | |
| <u>Melica californica</u> Scribn. | 1498 |
| Mixed conifer forest. Infrequent. | |
| <u>Sitanion hystrix</u> (Nutt.) J. G. Sm. | 1506 |
| Serpentine outcrops. Common. | |
| <u>Torreyochloa pauciflora</u> (Presl.) Church | 1480 |
| Seeps in mixed conifer forest. Common. | |
| <u>Vulpia</u> sp. | 1475 |
| Serpentine outcrops. Frequent. | |

HYDROPHYLLACEAE

| | |
|---------------------------------|-----|
| <u>Phacelia corymbosa</u> Heps. | 534 |
| Serpentine outcrops. Frequent. | |

IRIDACEAE

| | |
|--------------------------------------|------|
| <u>Iris tenuissima</u> Dykes | 1390 |
| Serpentine outcrops. Frequent. | |
| <u>Sisyrinchium idahoense</u> Bickn. | 1393 |
| Silver Spring. Frequent. | |

JUNCACEAE

| | |
|---|------|
| <u>Juncus covillei</u> Piper | 1407 |
| Swim Meadow; seeps. Frequent. | |
| <u>Juncus effusus</u> L. var. <u>gracilis</u> Hook. | 1404 |
| Swim Meadow; Silver Spring. Frequent. | |
| <u>Juncus ensifolius</u> Wikstr. | 1412 |
| Seeps. Frequent. | |

LABIATAE

| | |
|---|-----------|
| <u>Monardella odoratissima</u> Benth. ssp. <u>pallida</u> (Heller) Epl. | 713 |
| Serpentine outcrops; ephemeral streambeds. Frequent. | |
| <u>Prunella vulgaris</u> L. ssp. <u>lanceolata</u> (Barton) Hult. | 697 |
| Riparian; seeps. Common. | |
| <u>Pycnanthemum californicum</u> Torr. | 716 |
| Ephemeral streambeds. Frequent. | |
| <u>Scutellaria antirrhinoides</u> Benth. | 695 |
| Ephemeral streambeds; seeps. Common. | |
| <u>Stachys mexicana</u> Benth. | 698, 1028 |
| Riparian. Common. | |

LEGUMINOSAE

| | |
|--|-----|
| <u>Cercis occidentalis</u> Torr. ex Gray | 799 |
| Serpentine outcrops. Frequent. | |
| <u>Lathyrus polyphyllus</u> Nutt. ex T. & G. | 91 |
| Coniferous forests, Frequent. | |
| <u>Lotus grandiflorus</u> (Benth.) Green | 845 |
| Silver Spring. | |
| <u>Lotus purshianus</u> (Benth.) Clem. & Clem. | 847 |
| Mixed conifer forest. Frequent. | |
| <u>Lupinus latifolius</u> J.G. Agardh | 832 |
| Mixed conifer forest. Infrequent. | |
| <u>Lupinus stiversii</u> Kell. | 812 |
| Banks along Smoky Creek, Uncommon. | |
| <u>Trifolium monanthum</u> Gray | 839 |
| Meadows. Infrequent | |
| <u>Trifolium</u> sp. | |
| Along dry streambeds. Uncommon. | |

LILIACEAE

| | |
|---|------|
| <u>Disporum hookeri</u> (Torr.) Nichols | 1332 |
| var. <u>trachyandrum</u> (Torr.) Q. Jones | |
| Mixed conifer forest. Frequent. | |
| <u>Hastingsia alba</u> (Durand) S. Watson | 1329 |
| Swim Meadow; Silver Springs. Common. | |
| <u>Lilium pardalinum</u> Kell. | 1342 |
| Seeps. Frequent. | |
| <u>Smilacina racemosa</u> (L.) Desf. | 1331 |
| Rocky talus slopes. Common. | |

MALVACEAE

| | |
|------------------------------------|-----|
| <u>Sidalcea glaucescens</u> Greene | 134 |
| Swim Meadow. Uncommon. | |

ONAGRACEAE

| | |
|---|-----|
| <u>Epilobium adenocaulon</u> Hausskn. var. <u>parishii</u> (Trel.) Munz | 932 |
| Swim Meadow. Frequent. | |
| <u>Epilobium paniculatum</u> Nutt. ex T. & G. | 929 |
| Swim Meadow. Frequent. | |

ORCHIDACEAE

| | |
|---|------|
| <u>Epipactis gigantea</u> Dougl. ex Hook. | 1399 |
| Silver Spring. Frequent. | |
| <u>Habenaria dilata</u> (Pursh) Hook. | 1396 |
| var. <u>leucostachys</u> (Lindl.) Ames | |
| Silver Spring. Frequent. | |

PINACEAE

| | |
|--|----|
| <u>Abies concolor</u> (Gord. & Glend.) Lindl. | 49 |
| Mixed conifer forest; seeps. Infrequent. | |
| <u>Pinus jeffreyi</u> Grev. & Balf. in A. Murr. | 54 |
| Serpentine outcrops. Common. | |
| <u>Pinus lambertiana</u> Dougl. | 52 |
| Mixed conifer forest; rocky slopes. Common. | |
| <u>Pinus ponderosa</u> Dougl. ex P. & C. Lawson | 53 |
| Mixed conifer forest. Common. | |
| <u>Pinus sabiniana</u> Dougl. | 55 |
| Serpentine outcrops. Infrequent. | |
| <u>Pseudotsuga menziesii</u> (Mirb.) Franco | 57 |
| Mixed conifer forest; serpentine outcrops. Common. | |

POLYGALACEAE

| | |
|-----------------------------------|-----|
| <u>Polygala cornuta</u> Kell. | 156 |
| Mixed conifer forest. Infrequent. | |

POLYGONACEAE

| | |
|---|-----|
| <u>Eriogonum compositum</u> Dougl. ex Benth. | 45S |
| Serpentine outcrops. Infrequent. | |
| <u>Eriogonum nudum</u> Dougl. ex Benth. var. <u>oblongifolium</u> Wats. | 68S |
| Serpentine outcrops. Frequent. | |
| <u>Eriogonum strictum</u> Benth. | 65S |
| ssp. <u>proliferum</u> (T. & G.) S. Stokes | |
| Serpentine outcrops. Frequent. | |
| <u>Eriogonum</u> sp. | 33S |
| Serpentine outcrops. Infrequent. | |
| <u>Polygonum spargulariaeforme</u> Meissn. | 362 |
| Rocky slopes. Frequent. | |
| <u>Polygonum</u> sp. | 360 |
| Rocky slopes. Frequent. | |
| <u>Rumex californicus</u> Rech. | 357 |
| Seeps and vernal pond. Frequent. | |

PRIMULACEAE

| | |
|-----------------------------------|-----|
| <u>Dodecatheon</u> sp. | 400 |
| Grassy, rocky slopes. | |
| <u>Trientalis latifolia</u> Hook. | 404 |
| Mixed conifer forest. Common. | |

PYROLACEAE

| | |
|--|-----|
| <u>Chimaphila menziesii</u> (R. Br. ex D. Don) Spreng. | 435 |
| Mixed conifer forest. Infrequent. | |
| <u>Chimaphila umbellata</u> (L.) Barton | 435 |
| var. <u>occidentalis</u> (Rydb.) Blake | |
| Mixed conifer forest. Frequent. | |

PYROLACEAE

- Pyrola picta Sm. 434
Mixed conifer forest. Infrequent.
Pyrola picta Sm. forma aphylla (Sm.) Camp. 434
Mixed conifer forest. Rare.

RANUNCULACEAE

- Aquilegia formosa Fisch. in DC. 105
var. truncata (F. & M.) Baker
Riparian; seeps in mixed conifer forest. Common.
Clematis ligusticifolia Nutt. in T. & G. 103
Seeps. Infrequent.
Delphinium sp. 81
Riparian. Infrequent.

RHAMNACEAE

- Ceanothus cuneatus (Hook) Nutt. 982
Rocky, serpentine slopes. Common.
Ceanothus integerrimus H. & A. 977
Rocky slopes. Common.
Ceanothus prostratus Benth. 984
Mixed conifer forest. Infrequent.
Rhamnus californica Esch. 972
Serpentine outcrops Frequent.
Rhamnus purshiana DC. 973
Riparian; seeps. Frequent.

ROSACEAE

- Amelanchier pallida Greene 793
Riparian. Frequent.
Cercocarpus betuloides Nutt. ex T. & G. 782
Rocky, serpentine outcrops. Infrequent.
Potentilla glandulosa Lindl. 774
Ephemeral streambeds. Infrequent.
Rosa pisocarpa Gray 787
Riparian, seeps. Common.
Rubus parviflorus Nutt. 785
Riparian; seeps. Common.
Rubus ursinus Cham. & Schlecht. 784
Riparian; seeps. Common.

RUBIACEAE

- Galium ambiguum Wight var. siskiyouense Ferris 1041
Serpentine outcrops. Frequent.
Galium nuttallii Gray 1042
Serpentine outcrops. Frequent.
Galium triflorum Michx. 1040
Riparian; seep. Common.

SALICACEAE

- Salix laevigata Bebb. 913
Riparian. Frequent.

SAXIFRAGACEAE

- Heuchera micrantha Dougl. ex Lindl. 741
Rocky slopes. Uncommon.
Parnassia palustris L. var. californica Gray 731
Silver Spring. Frequent.
Philadelphus lewisii Pursh ssp. californicus (Benth.) Munz 744
Riparian. Frequent.
Whipplea modesta Torr. 745
Mixed conifer forest; seeps. Infrequent.

SCROPHULARIACEAE

- Mimulus guttatus Fisch. ex DC. 616
Swim Meadow; Silver Spring. Frequent.
Mimulus moschatus Dougl. ex Lindl. 610
Seeps. Common.
Penstemon corymbosus Benth. 641
Rocky slopes. Uncommon.
Veronica americana (Raf.) Schw. 656
Seeps. Frequent.

SINOPTERIDACEAE

- Cheilanthes gracillima D.C. Eat. in Torr. 34
Rocky slopes. Frequent.
Pellaea brachyptera (T. Moore) Baker 36
Serpentine outcrops. Frequent.

TAXACEAE

- Taxus brevifolia Nutt. 65
Riparian; seeps. Uncommon.

UMBELLIFERAE

- Ligusticum sp. 1013
Riparian. Frequent.
Lomatium sp. 1018
Rocky slopes. Frequent.
Perideridia gairdneri (H. & A.) Math. 1013
Swim Meadow. Infrequent.

VERBENACEAE

- Verbena hastata L. 687
Riparian. Infrequent.


VITACEAE

- Vitis californica Benth. 969
Riparian. Infrequent.

Voucher specimens collected on the proposed Smoky Creek Research Natural Area and vicinity. The collection numbers given are those of Dean Wm. Taylor. Specimens are deposited in the Herbarium, University of California, Davis. Those indicated by HSU are duplicates sent to Humboldt State University, Arcata.

- 7696A *Galium triflorum* Michx.
- 7696B *Juncus effusus* L. var. *gracilis* Hook.
- 7697 *Datisca glomerata* (Presl.) Baill.
- 7698 *Juncus covillei* Piper HSU
- 7699 *Agrostis exartha* Trin.
- 7700 *Carex exsiccata* Bailey
- 7701 *Monardella odoratissima* Benth.
- 7703 *Haplopappus ophitidis* (J. T. Howell) Keck
- 7704 *Wyethia angustifolia* (DC.) Nutt.
- 7705 *Helianthus exilis* Gray HSU
- 7706 *Pycnanthemum californicum* Torr. HSU
- 7707 *Scutellaria antirrhinoides* Benth.
- 7708 *Cuscuta subinclusa* Dur. & Hilg.
- 7709 *Polygonum spergulariaeforme* Meissn. HSU
- 7710 *Perideridia gairdneri* (H. & A.) Math.
- 7711 *Chrysothamnus parryi* (Gray) Greene ssp. *latior* Hall & Clem.
- 7714 *Lactuca saligna* L.
- 7715 *Hastingsia alba* (Durand) S. Wats.
- 7716 *Juncus ensifolius* Wikstr.
- 7717 *Cirsium breweri* (Gray) Jeps. HSU
- 7718 *Helenium bigelovii* Gray
- 7719 *Sidalcea glaucescens* Greene
- 7720 *Aster adscendens* Lindl.
- 7721 *Parnassia palustris* L. var. *californica* Gray
- 7722 *Epipactis gigantea* Dougl. ex Hook.
- 7723 *Rhynchospora glomerata* (L.) Vahl. var. *minor* Britton. HSU
- 7723 *Carex buxbaumii* Wahl.
- 7725 *Lotus purshianus* (Benth.) Clem. & Clem.

Map 1. Regional map showing the general location of the proposed Smoky Creek Research Natural Area.




Smoky Creek




Map 3.

Topographic map showing the location of the proposed Smoky Creek Research Natural Area. The solid line shows the boundaries as proposed by the Shasta-Trinity National Forest. Our proposed boundaries are shown by the dotted line. The upper boundary could be left at Forest Road 29 if it is felt that inclusion of additional Pinus jeffreyi Habitat types is not needed. (Dubakella 15' map)



(The dots plot the location of several rare plant taxa in the vicinity. See text for details. The taxa are coded by 2 letter abbreviations of the generic name and the specific epithet). ALHO: *Allium hoffmanii*; HAOP: *Haplopappus ophitidis*; ARRO: *Arenaria rosei*; SATR: *Sanicula tracyi*; ASSO: *Asclepias solanoana*.






Table 1. Climatic summary for the Forest Glen station nearby the proposed Smoky Creek Research Natural Area.

LOC: Forest Glen Ranger Station, Trinity County, California. 2343 feet elevation.

Means taken from 1973 California Annual Summary.

LAT: 40.22 N

MAX SS: 100 mm

| | J | F | M | A | M | J | J | A | S | O | N | D | YR |
|----|-----|-----|-----|-----|------|------|------|------|------|------|-----|-----|------|
| T | 1.9 | 4.2 | 6.6 | 9.8 | 14.8 | 16.3 | 20.2 | 19.9 | 16.6 | 11.3 | 5.7 | 2.8 | 10.8 |
| PE | 5 | 12 | 26 | 46 | 85 | 96 | 126 | 115 | 81 | 47 | 18 | 7 | 664 |
| P | 304 | 243 | 196 | 97 | 114 | 23 | 7 | 3 | 20 | 103 | 168 | 296 | 1572 |
| SS | 100 | 100 | 100 | 100 | 100 | 48 | 14 | 4 | 2 | 58 | 100 | 100 | |
| AE | 5 | 12 | 26 | 46 | 85 | 75 | 40 | 13 | 22 | 47 | 18 | 7 | 394 |
| - | | | | | | 21 | 86 | 103 | 60 | | | | 269 |
| + | 299 | 231 | 170 | 51 | 29 | | | | | 0 | 108 | 289 | 1177 |

KEY TO SYMBOLS

— PPT greater than scale

+ PPT

x Mean Monthly Temperature

o Mean Monthly Potential Evapotranspiration

. Soil Water storage/utilization




Table 2

| ALLIANCE | | Alnus rhombifolia-Peltiphyllum peltatum | | | | | | | | D. caespitosa | | Carex rostrata | | | |
|-----------------------------------|--|---|--------|--------|--------|--------|--------|--------|--------|-----------------------------------|--------|---|--------|-----------------|------|
| ASSOCIATION | | Alnus rhombifolia-Gallium triflorum | | | | | | | | Prunella vulgaris-Carex bolanderi | | Deschampsia caespitosa-Carex bauxbaumii | | Carex exsiccata | |
| Plot or Relevé Code | | S H | S H | S H | S H | S H | S H | S H | S H | S H | S H | S H | S H | S H | |
| | | 2 | 2 | P | 4 | 6 | P | 7 | 2 | 1 | 1 | 1 | 1 | 1 | |
| | | 3 | 4 | 3 | | | 1 | | 5 | 4 | 7 | 8 | 9 | | |
| ELEVATION(m) | | 1219 | 1127 | 994 | 1355 | 1274 | 1222 | 990 | 1359 | 1143 | 1219 | 1271 | 1030 | 1029 | 1027 |
| SLOPE | | 0° | 1° | 2° | 2° | 2° | 2° | 2° | 0° | 1° | 6° | 3° | 10° | 12° | 15° |
| ASPECT | | SE | N | SSE | NE | N | ESE | NE | WNW | W | N | SW | SE | SSE | SE |
| SUBSTRATE | | PCm | PCm | PCm | PCm | PCm | PCm | PCm | PCm | PCm | PCm | PCub | PCub | PCub | PCub |
| ALNUS RHOMBIFOLIA | | 2 | 1 | 3 | 1 | 2 | 4 | 1 | | | | | | | |
| ACER MACROPHYLLUM | | 2 | + | 1 | 1 | | | 1 | | | | | | | |
| ELYMUS GLAUCUS | | + | | + | 1 | 2 | 1 | 1 | + | + | | | | | |
| ROSA PISOCARPA | | | + | | | | + | + | + | | | | | | |
| GALIUM TRIFLORUM | | 1 | 1 | | 1 | 1 | 1 | 1 | | 2 | | | | | |
| CAREX NUDATA | | + | + | 2 | 2 | 2 | | | | | | | | | |
| ARALIA CALIFORNICA | | 1 | + | 2 | | + | | 1 | | | | | | | |
| PTERIDIUM AQUILINUM PUBESCENS | | 1 | | | | | + | 1 | | | | | | | |
| STACHYS MEXICANA | | | + | | | | 2 | 1 | | 2 | | | | | |
| AQUILEGIA FORMOSA TRUNCATA | | | + | + | 1 | | 1 | + | | | | | | | |
| DATISCA GLOMERATA | | | | | 1 | + | 1 | + | | | 1 | | | | |
| CHRYSOLEPSIS CHRYSOPHYLLA | | 1 | + | | | | 2 | | | | | | | | |
| PHILADELPHIS LEWISTII GORDONIANUS | | | | 1 | | + | | + | | | | | | | |
| SCUTILLARIA ANTHIRNOIDES | | + | | | 2 | 2 | | | | | | | | | |
| CAREX BOLANDERI | | | | | | | 1 | + | 1 | 1 | | | | | |
| TORREYOCHLOA PAUCIFLORA | | | | + | | | 3 | | | 2 | | | | | |
| CAREX AMPLIFOLIA | | | | | | | 1 | 1 | | 2 | | | | | |
| RUBUS LEUCODERMIS | | | | | | | 2 | + | + | | | | | | |
| EQUISETUM LAEVIGATUM | | | | | | | 1 | 1 | | 2 | | | | | |
| VERONICA AMERICANA | | | | | | | + | | | 1 | | | | | |
| CIRSIIUM ARVENSE | | | | | | | 1 | | + | + | | | | | |
| PRUNELLA VULGARIS LANCEOLATA | | | | | 1 | + | 1 | 1 | 3 | 1 | 1 | 1 | 1 | + | |
| DESCHAMPSIA CAESPITOSA | | | | | | | | | | | 3 | 2 | 2 | 2 | |
| JUNCUS COVILLEI | | | | | | | + | | 1 | | 1 | 1 | 2 | 2 | + |
| JUNCUS EFFUSUS GRACILIS | | | | | | | 3 | | | + | 1 | 3 | 1 | 1 | |
| CAREX BAUXBAUMII | | | | | | | | | | | 1 | 3 | 2 | 2 | |
| CIRSIIUM BREWEI | | | | | | | | | | | + | + | + | 1 | |
| HASTINGSIA ALBA | | | | | 2 | | | | | | + | 1 | + | 1 | |
| HELENIUM BIGELOVIT | | | | | | | | | | + | 1 | 1 | + | + | |
| MIMULUS GUTTATUS | | | | | | | | | | | 1 | + | + | + | |
| EQUISETUM ARVENSE | | | | | | | | | | + | | 1 | + | + | |
| EPILOBIUM ADENOCAULON | | | | | | | + | | | + | | | + | + | |
| PARNASSIA PALUSTRIS CALIFORNICA | | | | | | | | | | | | + | 2 | 2 | |
| JUNCUS ENSIFOLIUS | | | | | | | | | | | | 1 | + | 1 | |
| RHYNCHOSPORA GLOMERATA MINOR | | | | | | | | | | | | 1 | + | R | |
| EPIPACTIS GIGANTEA | | | | | | | | | | | | 1 | 1 | + | |
| SISYRINCHIUM IDAHOENSE | | | | | | | | | | | | 1 | 1 | + | |
| HABENARIA DILATATA LEUCOSTACHYS | | | | | | | | | | | | + | 1 | | |
| CAREX EXSICCATA | | | | | | | | | | | | | | | 5 |
| RUMEX CALIFORNICUS | | | | | | | | | 1 | | + | | | | + |
| MIMULUS MOSCHATUS | | | | | | | | | 1 | | | | | | |
| VITIS CALIFORNICA | | | | 1 | | | | | | | | | | | |
| PERIDERIS GARDNERI | | | | | | | | | | | + | | | | |
| SIDALCEA GLAUDESCENS | | | | | | | | | | | + | | | | |
| HELEOCHARIS PARTISHTI | | | | | | | | | | | 1 | | | | |
| ASTER ASCENDENS | | | | | | | | | | | 1 | | | | |
| POTENTILLA GLANDULOSA NEVADENSIS | | | | | | | + | | | | | | | | |
| PYCANTHEMUM CALIFORNICUM | | | | | | | 1 | | | | | | | + | |
| ARTEMISIA DOUGLASSII | | | | | 1 | 1 | | | | | | | | | |
| AGROSTIS EXARATA | | | | | | | | | | | | | | | + |
| TRIFOLIUM MONANTHUM | | | | | | | | | | + | | | | | |
| LILIUM PAROALINUM | | | | + | | | | | | | + | | | | |
| RUDDOWARIA FIMBRATA | | | | | | | | | | | | | | | |
| PETESITES PALMATA | | | | | | | | 2 | | | | | | | |
| LOTUS GRANDIFLORUS | | | | | + | | | | | 1 | | | | 1 | |
| ABIES CONCOLOR | | | | | | | | + | | | | | | | |
| AMELANCHIER PALLIDA | | | | | | | + | | | | | | | | |
| EUONYMUS OCCIDENTALIS | | | | | | | | | + | | | | | | |
| EUPATORIUM OCCIDENTALE | | | | + | | + | | | | | | | | | |

Table 3

[illegible][illegible]

HAPLOPAPPUS OPHETINUS
PHACELIA CORYMBOSA
ERIOPHYLLUM LANATUM LANCEOLATUM
ERIOGONUM STRICTUM
CIRSIO CALLIPES
SENECIO GREYNEI

QUERCUS KELLOGGII
CURYULUS CORNUA CALIFORNICA
HEIRACIUM ALBIFLORUM
CHINAPHILA UMBELLATA OCCIDENTALIS
BROMUS MARGINATUS
LATHYRUS POLYPHYLLUS
PYROLA PICTA
APUCYNUM PINULID

| | | | | | | | | | |
|---|---|---|---|---|---|--|---|---|---|
| | | 2 | 7 | 1 | * | | ? | 1 | 1 |
| R | 2 | 2 | | | * | | | | |
| | * | * | * | * | * | | | | |
| | * | * | * | * | * | | | | * |
| | | 1 | 1 | * | * | | * | | |
| | * | 1 | | | | | ? | | |
| | * | * | | | * | | * | | |
| | 1 | * | * | | | | | | |

[illegible]

| | | | |
|------------------------------------|---|---|-----|
| CARYLEA LANULOSA | | . | |
| CAREX GLOBOSA | | . | |
| MELLIANTHUS EXILIS | | . | |
| MEETHIA ANGUSTIFOLIA | | . | |
| BALSAMORHIZA SACITTATA | | . | . |
| GALIUM AMBIGUUM SISKIYOUENSE | + | . | . |
| PINUS SABINIANA | | 1 | . |
| CERCIS OCCIDENTALIS | 1 | 2 | |
| RHOMUS TECTORUM | | | |
| HADIA ELEGANS | | | |
| CHRYSOETHAMNUS PARRYI LATICE | | 1 | R |
| STREPIANTHUS TOPIONENSIS PALLIDUS | | | |
| ZAUSCHNERIA CALIFORMICA LATIFOLIA | | | |
| MUNARDELLA ODDRATISSIMA PALLIDA | | 1 | . |
| SYPHOEDRICARPUS VOLLIS | | | 1 . |
| CURMUS MUTTALLII | | | . |
| AMBUTUS MENZIESII | | | . |
| CAREX MULTICAULIS | | | 1 |
| IMIS TENUISSIMA | | . | . |
| PHURGUEHENDON FLAVESCENS VILLERSEN | | . | . |

Table 4.

Summary of forest plot data for the Smoky Creek area. Upper values are means for all plots of a given type, and lower values are Coefficient of Variation expressed as a percentage of the mean value.

| Taxon | Basal Area (m ² /ha) | Rel. Dom. | Dens- ity (Stems/ ha) | Rel. Dom. | Impt. Value | \bar{X} DBH (cm) |
|---|---------------------------------------|---------------|--------------------------------|---------------|----------------|--------------------------|
| A. <i>Alnus rhombifolia</i> - <i>Galium triflorum</i> Association. Plots 1,3. | | | | | | |
| <i>Alnus rhombifolia</i> | 11.0 (12) | 58.0 (63) | 111.3 (20) | 52.5 (74) | 110.5 (68) | 34.3 (01) |
| <i>Abies concolor</i> | 7.3 (141) | 23.1 (141) | 94.5 (141) | 25.0 (141) | 48.2 (141) | 12.0 (141) |
| <i>Acer macrophyllum</i> | 1.1 (141) | 7.8 (141) | 15.9 (141) | 10.0 (141) | 17.8 (141) | 15.0 (141) |
| <i>Pseudotsuga menziesii</i> | 3.1 (141) | 9.9 (141) | 15.9 (141) | 4.1 (141) | 14.0 (141) | 25.0 (141) |
| <i>Taxus brevifolia</i> | 0.2 (141) | 0.8 (141) | 15.9 (141) | 4.1 (141) | 5.0 (141) | 7.5 (141) |
| <i>Chrysolepis chrysophylla</i> | 0.1 (141) | 0.1 (141) | 15.9 (141) | 4.1 (141) | 4.1 (141) | 2.0 (141) |
| B. <i>Pinus jeffreyi</i> - <i>Haplopappus ophitidis</i> Association. Plots 7,8. | | | | | | |
| <i>Pinus jeffreyi</i> | 36.2 (64) | 87.5 (15) | 302.7 (37) | 65.0 (10) | 152.8 (13) | 35.7 (73) |
| <i>Calocedrus decurrens</i> | 3.4 (76) | 12.0 (107) | 175.0 (64) | 35.0 (20) | 47.1 (42) | 14.7 (63) |
| C. <i>Pinus jeffreyi</i> - <i>Calamagrostis koelerioides</i> Association. Plots 4,9,13. | | | | | | |
| <i>Pinus jeffreyi</i> | 32.8 (32) | 65.3 (45) | 381.9 (88) | 67.6 (53) | 132.2 (46) | 39.3 (57) |
| <i>Calocedrus decurrens</i> | 24.1 (98) | 43.8 (29) | 180.4 (90) | 16.2 (87) | 45.2 (88) | 25.8 (94) |
| <i>Pseudotsuga menziesii</i> | 3.7 (170) | 5.1 (170) | 190.9 (159) | 9.8 (146) | 21.1 (159) | 8.2 (11) |

Table 4 concluded.

| Taxon | BA | RD | D | RD | IV | \bar{X} |
|--|---------------|---------------|---------------|---------------|---------------|--------------|
| D. <i>Pseudotsuga menziesii</i> - <i>Pinus ponderosa</i> Association. Plots 2,5,6,11,12,14,15. | | | | | | |
| <i>Pseudotsuga menziesii</i> | 49.6 (61) | 53.7 (57) | 498.2 (63) | 48.7 (43) | 102.1 (45) | 24.4 (36) |
| <i>Pinus lambertiana</i> | 14.6 (125) | 21.7 (139) | 104.0 (80) | 11.7 (86) | 33.6 (117) | 27.0 (92) |
| <i>Pinus ponderosa</i> | 18.4 (186) | 19.1 (121) | 95.0 (131) | 11.3 (133) | 30.5 (113) | 37.0 (91) |
| <i>Quercus chrysolepis</i> | 2.2 (181) | 2.4 (168) | 157.5 (82) | 16.0 (84) | 17.9 (84) | 8.6 (65) |
| <i>Quercus kelloggii</i> | 1.1 (134) | 1.7 (143) | 62.6 (87) | 7.3 (90) | 9.0 (100) | 8.7 (83) |
| <i>Cornus nuttallii</i> | 0.4 (280) | 0.7 (280) | 9.0 (280) | 0.6 (280) | 1.3 (280) | 3.1 (280) |
| <i>Quercus garryana</i> | 0.1 (280) | 0.1 (280) | 9.0 (280) | 0.7 (280) | 0.9 (280) | 1.7 (280) |
| <i>Calocedrus decurrens</i> | 0.0 (280) | 0.0 (280) | 6.0 (280) | 0.6 (280) | 0.6 (280) | 8.6 (280) |

Summary Table of Forest Statistics for the proposed Smoky Creek Research Natural Area. Values given are for all plots sampled, and are means and Coefficient of Variation expressed as a percentage of the mean.

| Taxon | Basal Area (m ² /ha) | Rel. Dom. | Dens- ity (Stems/ ha) | Rel. Dens. | Impt. Val. | \bar{X} |
|---------------------------------|---------------------------------------|---------------|--------------------------------|---------------|---------------|---------------|
| <i>Pseudotsuga menziesii</i> | 24.4 (130) | 52.1 (216) | 273.3 (120) | 26.5 (105) | 53.9 (107) | 15.8 (100) |
| <i>Pinus jeffreyi</i> | 11.4 (159) | 25.1 (153) | 118.1 (190) | 25.5 (173) | 50.7 (141) | 13.0 (163) |
| <i>Calocedrus decurrens</i> | 5.3 (251) | 7.4 (215) | 63.2 (172) | 8.2 (169) | 16.7 (162) | 7.5 (188) |
| <i>Pinus lambertiana</i> | 6.8 (206) | 10.3 (223) | 48.5 (158) | 15.4 (164) | 15.7 (198) | 12.6 (169) |
| <i>Alnus rhombifolia</i> | 1.4 (265) | 7.7 (298) | 14.9 (266) | 7.0 (302) | 14.7 (298) | 4.6 (262) |
| <i>Pinus ponderosa</i> | 9.3 (268) | 8.9 (203) | 44.3 (215) | 5.2 (217) | 14.2 (193) | 18.3 (158) |
| <i>Quercus garryana</i> | 0.8 (357) | 6.3 (383) | 12.7 (280) | 3.6 (350) | 10.0 (371) | 2.4 (282) |
| <i>Quercus chrysolepis</i> | 1.0 (276) | 1.0 (254) | 73.6 (160) | 7.4 (162) | 8.5 (164) | 4.0 (144) |
| <i>Abies concolor</i> | 1.0 (366) | 3.1 (380) | 27.5 (264) | 4.5 (294) | 7.7 (325) | 2.0 (292) |
| <i>Quercus kelloggii</i> | 0.5 (216) | 0.8 (230) | 29.2 (166) | 3.4 (168) | 4.2 (174) | 4.0 (161) |
| <i>Taxus brevifolia</i> | 0.1 (392) | 0.1 (387) | 2.2 (387) | 0.5 (387) | 0.6 (387) | 1.0 (387) |
| <i>Acer macrophyllum</i> | 0.1 (387) | 1.0 (387) | 2.1 (387) | 1.3 (387) | 2.3 (387) | 2.0 (387) |
| <i>Cornus nuttallii</i> | 0.2 (387) | 0.3 (387) | 4.2 (387) | 0.3 (387) | 0.6 (387) | 1.4 (387) |
| <i>Chrysolepis chrysophylla</i> | 0.0 (387) | 0.0 (387) | 2.1 (387) | 0.5 (387) | 0.5 (387) | 0.2 (387) |

BASAL AREA PROGRAM
STAND No.-5MKP5
PLOT RADIUS= 15.000 m
PLOT AREA = 706.85 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|------------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pseudotsuga menziesii</i> | 46.230 | 63.69 | 565.88 | 60.6 | 124.3 | 18.5 |
| <i>Pinus ponderosa</i> | 14.756 | 20.33 | 28.29 | 3.0 | 23.3 | 81.2 |
| <i>Pinus lambertiana</i> | 9.751 | 13.43 | 28.29 | 3.0 | 16.4 | 58.0 |
| <i>Quercus chrysolepis</i> | 1.236 | 1.70 | 212.20 | 22.7 | 24.4 | 6.7 |
| <i>Quercus kelloggii</i> | .449 | .61 | 56.58 | 6.0 | 6.6 | 9.4 |
| <i>Calocedrus decurrens</i> | .155 | .21 | 42.44 | 4.5 | 4.7 | 5.3 |
| TOTALS | 72.580 | | 933.70 | | | |

BASAL AREA PROGRAM
STAND No.-5MKP6
PLOT RADIUS= 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|------------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pseudotsuga menziesii</i> | 69.440 | 41.92 | 318.30 | 38.4 | 80.3 | 31.0 |
| <i>Pinus lambertiana</i> | .022 | .01 | 31.83 | 3.8 | 3.8 | 3.0 |
| <i>Pinus ponderosa</i> | 95.319 | 57.54 | 318.30 | 38.4 | 96.0 | 47.6 |
| <i>Quercus chrysolepis</i> | .387 | .23 | 95.49 | 11.5 | 11.7 | 7.1 |
| <i>Quercus kelloggii</i> | .463 | .27 | 63.66 | 7.6 | 7.9 | 9.6 |
| TOTALS | 165.631 | | 827.60 | | | |

BASAL AREA PROGRAM
STAND No.-5MKP7
PLOT RADIUS= 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|-----------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pinus jeffreyi</i> | 19.832 | 78.71 | 381.97 | 60.0 | 138.7 | 17.2 |
| <i>Calocedrus decurrens</i> | 5.361 | 21.28 | 254.64 | 40.0 | 61.2 | 15.5 |
| TOTALS | 25.194 | | 636.61 | | | |

BASAL AREA PROGRAM
STAND No.-5MKP8
PLOT RADIUS= 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|-----------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pinus jeffreyi</i> | 52.764 | 96.98 | 222.81 | 70.0 | 166.9 | 54.3 |
| <i>Calocedrus decurrens</i> | 1.642 | 3.01 | 95.49 | 30.0 | 33.0 | 14.3 |
| TOTALS | 54.406 | | 318.30 | | | |

BASAL AREA PROGRAM

STAND No. = SNKP13

PLOT RADIUS = 10.000 meters

PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | T DB |
|-----------------------|----------------------------|--------------|-----------------|---------------|--------------|---------|
| Pinus jeffreyi | 41.740 | 46.70 | 732.11 | 74.1 | 120.9 | 20.8 |
| Calocedrus decurrens | 47.482 | 53.13 | 222.81 | 22.5 | 75.7 | 48.1 |
| Pseudotsuga menziesii | .145 | .16 | 31.83 | 3.2 | 3.3 | 7.6 |
| TOTALS | 89.368 | | 986.76 | | | |

BASAL AREA PROGRAM

STAND No. = SNKP14

PLOT RADIUS = 10.000 meters

PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | T DB |
|-----------------------|----------------------------|--------------|-----------------|---------------|--------------|---------|
| Pseudotsuga menziesii | 61.809 | 93.34 | 1145.91 | 85.7 | 179.0 | 22. |
| Cornus nuttallii | 3.292 | 4.97 | 63.66 | 4.7 | 9.7 | 22. |
| Quercus chrysolepis | 1.117 | 1.68 | 127.32 | 9.5 | 11.2 | 10. |
| TOTALS | 66.219 | | 1336.90 | | | |

BASAL AREA PROGRAM

STAND No. = SNKP15

PLOT RADIUS = 10.000 meters

PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | T DB |
|-----------------------|----------------------------|--------------|-----------------|---------------|--------------|---------|
| Pinus lambertiana | 38.078 | 39.35 | 159.15 | 20.0 | 59.3 | 50. |
| Pinus ponderosa | 4.939 | 5.10 | 31.83 | 4.0 | 9.1 | 44. |
| Pseudotsuga menziesii | 51.693 | 53.42 | 254.64 | 32.0 | 85.4 | 36. |
| Quercus chrysolepis | 1.509 | 1.56 | 286.47 | 36.0 | 37.5 | 7. |
| Quercus kelloggii | .529 | .54 | 63.66 | 8.0 | 8.5 | 10. |
| TOTALS | 96.750 | | 795.77 | | | |

BASAL AREA PROGRAM
STAND No.-SMKP1
PLOT RADIUS= 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | Y DBH |
|--------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| Alnus oregana | 10.074 | 31.88 | 95.49 | 25.0 | 56.8 | 34.6 |
| Taxus brevifolia | .562 | 1.77 | 31.83 | 8.3 | 10.1 | 15.0 |
| Pseudotsuga menziesii | 6.249 | 19.77 | 31.83 | 8.3 | 28.1 | 50.0 |
| Abies concolor | 14.674 | 46.43 | 190.98 | 50.0 | 96.4 | 24.0 |
| Castanopsis chrysophylla | .039 | .12 | 31.83 | 8.3 | 8.4 | 4.0 |
| TOTALS | 31.601 | | 381.97 | | | |

BASAL AREA PROGRAM
STAND No.-SMKP2
PLOT RADIUS= 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | Y DBH |
|-----------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| Pseudotsuga menziesii | 94.584 | 87.13 | 445.63 | 36.8 | 123.9 | 32.5 |
| Abies concolor | .832 | .76 | 222.81 | 18.4 | 19.1 | 6.4 |
| Quercus chrysolepis | 11.624 | 10.70 | 350.14 | 28.9 | 39.6 | 18.9 |
| Quercus garryana | .964 | .88 | 63.66 | 5.2 | 6.1 | 12.0 |
| Pinus lambertiana | .542 | .49 | 127.32 | 10.5 | 11.0 | 5.2 |
| TOTALS | 108.549 | | 1209.57 | | | |

BASAL AREA PROGRAM
STAND No.-SMKP3
PLOT RADIUS= 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | Y DBH |
|-------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| Alnus oregana | 12.014 | 84.22 | 127.32 | 80.0 | 164.2 | 34.0 |
| Acer macrophyllum | 2.249 | 15.77 | 31.83 | 20.0 | 35.7 | 30.0 |
| TOTALS | 14.264 | | 159.15 | | | |

BASAL AREA PROGRAM
STAND No.-SMKP4
PLOT RADIUS= 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | Y DBH |
|----------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| Pinus jeffreyi | 20.862 | 100.00 | 63.66 | 100.0 | 200.0 | 64.5 |
| TOTALS | 20.862 | | 63.66 | | | |

BASAL AREA PROGRAM
STAND No. = SMKP9
PLOT RADIUS = 10.000 m
PLOT AREA = 314.15 meters²

39B

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|------------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pinus jeffreyi</i> | 36.077 | 49.99 | 350.14 | 20.9 | 78.9 | 32.6 |
| <i>Pseudotsuga menziesii</i> | 11.061 | 15.32 | 541.12 | 44.7 | 60.0 | 8.9 |
| <i>Calocedrus decurrens</i> | 25.028 | 34.68 | 318.30 | 26.3 | 60.9 | 29.5 |
| TOTALS | 72.166 | | 1209.57 | | | |

BASAL AREA PROGRAM
STAND No. = SMKP10
PLOT RADIUS = 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|-------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pinus jeffreyi</i> | .665 | 5.38 | 127.32 | 50.0 | 55.3 | 6.5 |
| <i>Quercus garryana</i> | 11.691 | 94.61 | 127.32 | 50.0 | 144.6 | 25.0 |
| TOTALS | 12.356 | | 254.64 | | | |

BASAL AREA PROGRAM
STAND No. = SMKP11
PLOT RADIUS = 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|------------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pinus lambertiana</i> | 43.194 | 83.66 | 222.81 | 26.9 | 110.5 | 47.2 |
| <i>Pseudotsuga menziesii</i> | 2.610 | 5.05 | 222.81 | 26.9 | 31.9 | 11.3 |
| <i>Pinus ponderosa</i> | 3.204 | 6.20 | 222.81 | 26.9 | 33.1 | 11.4 |
| <i>Quercus kelloggii</i> | 2.613 | 5.07 | 159.15 | 19.2 | 24.3 | 10.5 |
| TOTALS | 51.628 | | 827.60 | | | |

BASAL AREA PROGRAM
STAND No. = SMKP12
PLOT RADIUS = 10.000 m
PLOT AREA = 314.15 meters²

| TAXON | BA (m ² /ha) | REL. DOM. | DENS. (#/ha) | REL. DENS. | IMP. VAL. | X DBH |
|------------------------------|----------------------------|--------------|-----------------|---------------|--------------|----------|
| <i>Pinus ponderosa</i> | 30.791 | 45.27 | 63.66 | 7.1 | 52.4 | 74.4 |
| <i>Pinus lambertiana</i> | 11.575 | 17.01 | 159.15 | 17.8 | 34.8 | 25.7 |
| <i>Pseudotsuga menziesii</i> | 21.552 | 31.68 | 541.12 | 60.7 | 92.4 | 18.9 |
| <i>Quercus chrysolepis</i> | .258 | .37 | 31.83 | 3.5 | 3.9 | 10.1 |
| <i>Quercus kelloggii</i> | 3.836 | 5.64 | 95.49 | 10.7 | 16.3 | 21.4 |
| TOTALS | 68.014 | | 891.26 | | | |

Appendix 2.

Typification of the vegetation associations sampled on the proposed Smoky Creek Research Natural Area. The Code of Phytosociological Nomenclature requires that vegetation syntaxa described in the literature for the first time after 1 January 1979 follow the rules set forth therein (cf. Vegetatio Vol. 32:131-185.). Unlatinized names for the various vegetation types can be referenced in Tables 2 & 3.

1. ALLIANCE *Peltiphylllo-Alnuetalia rhombifolii* All. Nov. Taylor 1979

Nomenclatural Type: *Galio-Alnetum rhombifolii* Taylor 1979

Alliance Diagnostic Taxa: *Alnus rhombifolia*, *Aralia californica*,
Stachys mexicana, *Peltiphyllum peltatum*,
Carex nudata.

1. ASSOCIATION *Galio-Alnetum rhombifolii* Assoc. Nov. Taylor 1979

Nomenclatural Type: Table 2, Relève SMK 23

Association Diagnostic Taxa: *Scutellaria anthirrnoides*, *Datisca glomerata*, *Carex nudata*, *Aralia californica*, *Galium triflorum*.

2. ASSOCIATION *Prunello-Caricetum bolanderi* Assoc. Nov. Taylor 1979

Nomenclatural Type: Table 2, Relève SMK 15

Association Diagnostic Taxa: *Carex bolanderi*, *Cirsium arvense*,
Torreychloa pauciflora, *Carex amplifolia*

2. ALLIANCE *Cirsio-Deschampsietalia* All. Nov. Taylor 1979

Nomenclatural Type: *Deschampsio-Caricetum bauxbaumii* Taylor 1979

Alliance Diagnostic Taxa: *Carex bauxbaumii*, *Deschampsia caespitosa*,
Juncus covillei, *Cirsium breweri*, *Hastingsia alba*, *Sisyrinchium idahoense*.

3. ASSOCIATION *Deschampsio-Caricetum bauxbaumii* Assoc. Nov. Taylor 1979

Nomenclatural Type: Table 2, Relève SMK 18

Association Diagnostic Taxa: *Cirsio-Deschampsietalia* taxa, *Rhynchospora glomerata* var. *minor*.

3. ALLIANCE *Caricon simulato-vesicariae* Beguin & Major 1975 em. Taylor 1979

Nomenclatural Type: Beguin and Major 1975, Table 3, Relève 7
(lectotypus Taylor 1979)

Alliance Diagnostic Taxa: *Carex simulata*, *C. vesicaria*.

4. ASSOCIATION *Caricetum exsiccatae* Assoc. Nov. Taylor 1979

Nomenclatural Type: Table, Relève SMK 1

Association Diagnostic Taxa: *Carex exsiccata*.

4. ALLIANCE Onychio-Pinion jeffreyi All. Nov. Taylor 1979

Nomenclatural Type: Haplopappo-Pinetum jeffreyi Taylor 1979

Alliance Diagnostic Taxa: Calamagrostis koelerioides, Onychium densum,
Rhamnus californica var, tomentella.

5. ASSOCIATION Haplopappo-Pinetum jeffreyi Assoc. Nov. Taylor 1979

Nomenclatural Type: Table 3, Releve SMK 8

Association Diagnostic Taxa: Haplopappus ophetidus, Phacelia
corymbosa, Cirsium callipes, Senecio
greenei, Arenaria rosei, Allium hoffmanii

6. ASSOCIATION Sitanio-Ceanothetum cuneatae Assoc. Nov. Taylor 1979

Nomenclatural Type: Table 3, Releve SMK

Association Diagnostic Taxa: Ceanothus cuneatus, Calamagrostis
koelerioides, Polygonum spargulariaefolium.

7. ASSOCIATION Calamagrostio-Pinetum jeffreyi Assoc. Nov. Taylor 1979

Nomenclatural Type: Table 3, Releve SMK 16

Association Diagnostic Taxa: Pinus jeffreyi, Calamagrostis
koelerioides, Quercus durata, Ceanothus
cuneatus.

5. ALLIANCE Pseudotsugion menziesii All. Nov. Taylor 1979

Nomenclatural Type: Pinio-Pseudotsugetum menziesii Taylor 1979

Alliance Diagnostic Taxa: Pseudotsuga menziesii, Pinus ponderosa,
Quercus chrysolepis, Pinus lambertiana,
Toxicodendron diversilobum.

8. ASSOCIATION Pinio-Pseudotsugetum menziesii Assoc. Nov. Taylor 1979

Nomenclatural Type: Table 3, Releve SMK 4

Association Diagnostic Taxa: Pinus ponderosa, Pseudotsuga menziesii.

Photo 1. Typical view of open to patchy dense *Pseudotsuga menziesii* *Pinus ponderosa* forest on the proposed Smoky Creek RNA.



Photo 2. View of the canopy of *Pseudotsuga*-*Pinus* forest, showing the characteristic structure of a mosaic of old-growth and younger successional stage stands.



Photo 3. View of typical *Pinus jeffreyi* vegetation, with scattered understory shrubs, all on serpentine.



Photo 4. *Ceanothus cuneatus* vegetation on rocky, serpentine sites with poor soil development.



Photo 5. Seep on serpentine at Silver Spring adjacent to the proposed Smoky RNA. Such vegetation is quite distinctive and of limited distribution in the region.

